

# THE MARVELLOUS PROOF OF THE GOLDBACH'S CONJECTURE

LESZEK W. GULA

*I dedicate my work to the city Lublin*

ABSTRACT. The proper proof of the Goldbach's Conjecture

## I. INTRODUCTION

On 7 June 1742, the German mathematician Christian Goldbach wrote a letter to Leonhard Euler in which he proposed the following conjecture: 'Every integer which can be written as the sum of two primes, can also be written as the sum of as many primes as one wishes, until all terms are units'. Euler replied in a last letter: 'Every even integer greater than 2 can be written as the sum of two primes, which is, thus, also a conjecture of Goldbach'. [1]

## II. THE GLDBACH'S CONJECTURE

Let

$$\mathbb{A} = \{3, 5, 7, \dots\} \wedge \mathbb{P} = \mathbb{A} \setminus \mathbb{A}_o \wedge \mathbb{P} \cap \mathbb{A}_o = \emptyset \wedge$$

$$\mathbb{A}_o = \{(2a + b) b : b \in \{3, 5, 7, \dots\} \wedge a \in \mathbb{N}\} = \{9, 15, 21, 25, 27, \dots\}.$$

**Conjecture 1** (Goldbach Conjecture). *For all  $g \in \{6, 8, 10, \dots\}$  and for some  $X, Y \in \mathbb{P}$ :*

$$g = X + Y.$$

*This is the conjecture.*

*Proof.*

$$\begin{aligned} \{6, 8, 10, \dots\} &= \{6, 12, 18, 24, 30, 36, 42, 48, 54, 60, 66, 72, 78, 84, 90, 96, \dots\} \cup \\ &\quad \{8, 14, 20, 26, 32, 38, 44, 50, 56, 62, 68, 74, 80, 86, 92, 98, \dots\} \cup \\ &\quad \{10, 16, 22, 28, 34, 40, 46, 52, 58, 64, 70, 76, 82, 88, 94, 100, \dots\} \Rightarrow \\ \mathbb{A}_1 &= \{3\} \wedge \mathbb{A}_2 = \{9, 15, 21, 27, 33, 39, 45, 51, 57, 63, 69, 75, 81, 87, 93, \dots\} \wedge \\ \mathbb{A}_3 &= \{7, 13, 19, 25, 31, 37, 43, 49, 55, 61, 67, 73, 79, 85, 91, 97, \dots\} \wedge \\ \mathbb{A}_4 &= \{5, 11, 17, 23, 29, 35, 41, 47, 53, 59, 65, 71, 77, 83, 89, 95, \dots\} \wedge \\ \mathbb{A}_3 \setminus \mathbb{P} &= \mathbb{A}_5 \wedge \mathbb{A}_4 \setminus \mathbb{P} = \mathbb{A}_6 \wedge \mathbb{A}_3 \setminus \mathbb{A}_o = \mathbb{A}_7 \wedge \mathbb{A}_4 \setminus \mathbb{A}_o = \mathbb{A}_8. \end{aligned}$$

---

*Date:* 19.05.2014 – 24.05.2014.

*1991 Mathematics Subject Classification.* Primary: 11P32; Secondary: 11D85.

*Key words and phrases.* Algebra of sets.

$$\begin{aligned} \{42, 48, 54, \dots\} &= \{g : g = x + y \wedge x, y \in \mathbb{A}_2\} \wedge \\ \{40, 46, 52, \dots\} &= \{g : g = x + y \wedge x \in \mathbb{A}_2 \wedge y \in \mathbb{A}_5\} \wedge \\ \{44, 50, 56, \dots\} &= \{g : g = x + y \wedge x \in \mathbb{A}_2 \wedge y \in \mathbb{A}_6\}. \end{aligned}$$

Hence for all  $g \in \{40, 42, 44, \dots\}$  and for some  $x, y \in \mathbb{A}_o$  :

$$g = x + y. \quad \blackbox$$

$$\{6\} = \{g : 3 + 3 \wedge 3 \in \mathbb{A}_1\}.$$

$$\{8\} = \{g : 3 + 5 \wedge 3 \in \mathbb{A}_1 \wedge 5 \in \mathbb{A}_8\}.$$

$$\begin{aligned} \{14, 20, 26, \dots\} &= \{g : g = X + Y \wedge X, Y \in \mathbb{A}_7\} \wedge \\ \{10, 16, 22, \dots\} &= \{g : g = X + Y \wedge X, Y \in \mathbb{A}_8\} \wedge \\ \{12, 18, 24, \dots\} &= \{g : g = X + Y \wedge X \in \mathbb{A}_7 \wedge Y \in \mathbb{A}_8\}. \end{aligned}$$

Hence for all  $g \in \{6, 8, 10, \dots\}$  and for some  $X, Y \in \mathbb{P}$  :

$$g = X + Y.$$

This is the proof. □

**Corollary 1.** For all  $g \in \{40, 42, 44, \dots\}$  and for some  $X, Y \in \mathbb{P}$  and for some  $x, y \in \mathbb{A}_o$  :

$$g = X + Y = x + y.$$

*This is the corollary.*

#### REFERENCES

- [1] [http://en.wikipedia.org/wiki/Goldbach's\\_conjecture](http://en.wikipedia.org/wiki/Goldbach's_conjecture)

LUBLIN-POLAND

E-mail address: lwgula@wp.pl